

LA-UR--86-4012

DE87 002895

TITLE: SPIN QUENCHING IN THE SO_8 AND SP_6 FERMION
MODELS OF COLLECTIVE MOTION

AUTHOR(S): Joseph N(atale) Gineocchio, T-5

SUBMITTED TO: Proceedings of the "XV International Colloquium
on Group Theoretical Methods in Physics", Drexel
University, Philadelphia, PA, October 20-24, 1986

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U S Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U S Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U S Department of Energy

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

MASTER

SPIN QUENCHING AND THE SO_8 AND SP_6 FERMION
MODELS OF COLLECTIVE MOTION

Joseph N. Ginocchio

Theoretical Division, Los Alamos National Laboratory
Los Alamos, NM 87545

ABSTRACT

The expectation value of the spin operator in the SO_8 and SP_6 models is studied.

1. INTRODUCTION

The SO_8 and SP_6 models¹⁾ assume that the low-lying collective states of heavy nuclei are dominated by collective pairs of fermions coupled to angular momentum $J^\pi = 0^+, 2^+$. These models are generalizations of the pairing model (SP_2) to include the quadrupole degree of freedom. Within these models, the total shell model space can be classified in terms of the number of nucleons not coupled to angular momentum zero or two. This classification follows from partitioning the single-particle angular momentum \vec{j} into pseudo-orbital angular momentum \vec{k} and pseudo-spin \vec{i} . For shell model orbitals for which $k = 1$ the generators of SP_6 are the multipole and pair operators that have pseudo-spin coupled to zero. For shell model orbitals with $i = 3/2$ the generators of SO_8 are the multipole and pair operators that have pseudo-orbital angular momentum coupled to zero. In this paper we focus on the SO_8 model.

2. SPIN IN THE COLLECTIVE SUBSPACE

The spin \vec{S} operator can be expanded in the $k-i$ basis from the $|j-m\rangle$ basis,

$$\vec{S} = \sum_{K,I} C(K,I) (b_{ki}^\dagger \bar{b}_{ki})^{(K,I)} \quad (1)$$

where $b_{km,ln}^\dagger$ creates a particle in the $k-i$ orbit with psuedo-orbital angular momentum projection m and psuedo-spin projection n , $()^{(K,I)}$ means coupled to total psuedo-orbital angular momentum K and total psuedo-spin I , and the coefficients $C(K,I)$ are given by,

$$C(K,I) = \sum_{j,j'} \frac{\langle j || \vec{S} || j' \rangle}{\sqrt{3}} \hat{j}' \hat{j} \hat{K} \hat{I} \left\{ \begin{matrix} kk & K \\ ii & I \\ jj' & 1 \end{matrix} \right\} . \quad (2)$$

where $\hat{j} = \sqrt{2j+1}$. We can rewrite this expression for the spin operator as

$$\vec{S} = q \vec{I} + \sum_{I,K \neq 0} C(K,I) (b^\dagger \bar{b})^{(K,I)} \quad (3)$$

where q is the quenching factor and is given by

$$q = \sqrt{(2/5\Omega)} C(K=0, I=1) . \quad (4)$$

For the SO_8 model the collective subspace made up only of $J^\pi = 0^+, 2^+$ pairs only; i.e., $u=0$, has $K=0$. Hence within this subspace only the first term in (4) contributes, and furthermore $J=I$. Thus the expectation value of the spin in the collective $u=0$ subspace depends only on q . We can derive a simple expression for q for heavy nuclei (i.e., the maximum angular momentum j is $l - 1/2$ where l is the real angular momentum). We get

$$q = \frac{1}{15} \left[-1 + \left[\frac{4}{\Omega} \right]^2 \pm 2 \sqrt{(1 - \left[\frac{2}{\Omega} \right]^2) (1 - 2 \left[\frac{2}{\Omega} \right]^2)} \right] \quad (5)$$

where $\Omega = \sum (2j+1)$ is one-half the shell degeneracy. The two signs arise from phase freedom in going from $k-i$ (or $i-k$ coupling) to the

j-j basis. Since the SO_8 algebra is invariant to unitary transformations, we can change the sign of the single-particle basis and leave the algebra invariant. This doesn't effect any diagonal matrix elements in the j-j basis, only off-diagonal matrix elements. The resulting q factors are given in Table 1.

Ω	q_+	q_-	q_j				
			j	<u>1</u>	<u>3</u>	<u>5</u>	<u>7</u>
				2	2	2	2
6	0.074	- 0.148	-0.333, 0.333, -0.143				
10	0.069	- 0.181	0.500	-0.200,	0.200,	-0.111	
∞	0.067	- 0.200					

Table 1. The q - factors vs. shell degeneracy for the SO_8 model in the first two columns and for the single-particle orbits in the remaining columns.

Hence we see that the absolute value of this factor is reduced in the SO_8 $u = 0$ subspace compared to the single-particle q-factors for the + phase convention. For the - phase convention it is reduced except for the orbit with the largest j. The amount of reduction depends on the phase convention.

3. CONCLUSION

We have shown that within the SO_8 model the expectation value of the spin operator in the subspace consisting only of $J^\pi = 0^+, 2^+$ pairs

4. REFERENCES

1. Ginocchio, J. N., Ann. of Physics 126, 234 (1980).
2. Wu, C.L., Feng, D.H., Chen, X.G., Chen, J.Q. and Guidry, M. W.,
Phys. Lett. B168, 313 (1986).